

# GenTORE

***Genomic management Tools to Optimise Resilience  
and Efficiency***

Grant agreement n°: 727213

**H2020 - Research and Innovation Action**

**Deliverable 6.2**

***Paper on identification of key weather perturbations in performance and effect of decrease in temperature***

**Due date:** M54 (Nov 2021)

**Actual submission date:** M54 (Nov 2021)

**Project start date:** 1<sup>st</sup> June 2017      **Duration:** 60 months

**Workpackage concerned:** WP6

**Concerned workpackage leader:** Eileen Wall

**Lead Beneficiary:** SRUC

**Dissemination level:**

- X PU:** Public (must be available on the website)  
 **CO:** Confidential, only for members of the consortium (including the Commission Services)  
 **CI:** Classified, as referred to in Commission Decision 2001/844/EC

## Table of content

1.	Summary.....	3
2.	Introduction .....	3
3.	Results .....	3
3.1	Average Weather.....	4
3.2	Extreme Weather.....	4
4.	Conclusions.....	5
5.	Partners involved in the work .....	5
6.	Annex – Paper .....	5

## 1. Summary

In this paper, submitted to animal, we investigated how beef cattle traits are affected by varying weather and frequency of extreme events. We analysed the effect of average daily precipitation and average daily maximum and minimum temperatures on a range of important carcass traits, including age at slaughter, cold carcass weight, carcass growth rate and conformation and fat score ( $N = >1.6$  million), as well as calf 200-day live weight and growth rate ( $N = >270\,000$ ), using data from abattoirs across Scotland, England and Wales and from calves in herds across Scotland. Animals which experienced higher daily maximum and lower daily minimum temperatures had poorer age at slaughter and carcass growth rates and calves had poorer 200-day weights. Increased precipitation also led to poorer cold carcass weights, conformation scores, calf 200-day weights and calf growth.

We also analysed the effect of frequency of extreme weather events, including heatwaves, cold waves, and dry and wet days. The frequency of heatwaves, dry and wet days were shown to have significant negative effects on almost all traits considered, predicting that an increase in frequency of heatwaves by 1 day per 100 days of life would reduce cold carcass weights by about 200g and increase age at slaughter by about 3 days.

Results show that varying weather and frequency of extreme weather, across the lifetime of a beef animal, influences traits which affect the potential profit for a beef farmer. These effects may be due to several factors, including direct effects on the animals resilience and efficiency, as well as feed availability and management decisions made by the farmer. However, there is potential to mitigate negative effects through a range of strategies.

## 2. Introduction

To identify climate resilient beef cattle, a key aim of GenTORE, we first need a better understanding of how they are affected by weather. We have good data on how cattle in the tropics are negatively affected by heat stress for a range of traits including milk yield, health and fertility and growth rates and some good evidence of dairy cattle in Europe experiencing negative effects of both extreme high and low temperatures. However, we lack large scale studies on the effects of weather on beef cattle in temperate environments. Beef cattle have higher upper critical temperatures than dairy cattle, so may be less affected by heat stress. However typically UK beef cattle are not housed as much as dairy which may mean they are more greatly affected by weather. Therefore, our aim was to investigate the effect of weather and frequency of extreme weather events on a range of cattle traits important to beef production in the UK.

## 3. Results

### 3.1 Average Weather

Almost all average weather parameters had a significant ( $p<0.05$ ) effect on every trait assessed (Table 1). In summary, increases in average daily maximum temperatures experienced by an animal resulted in poorer age at slaughter, carcass growth rate, calf 200-day weight and calf growth rate, but improved conformation and higher fat scores. Animals which experienced high average daily minimum temperature were associated with improved age at slaughter, cold carcass weight, conformation and fat scores, as well as higher calf 200-day weights, but poorer carcass and calf growth rates. Finally, animals which experienced greater average daily precipitation tended to be younger at slaughter and improved carcass growth rates, but poorer cold carcass weights, conformation and fat scores and poorer calf 200-day weights and calf growth rates.

**Table 1**

Table of solutions and standard errors (in brackets) for weather variables, including average daily maximum temperature (Tmax), average daily minimum temperature (Tmin) and average daily precipitation (Rain) and interactions from model for each carcass trait and calf trait. All effects are significant where given ( $p<0.05$ ). Non-significant effects are denoted by ns.

	Tmax	Tmin	Rain	Tmin x Tmax	Tmin x Rain
<b>Age at Slaughter (days)</b>	10.17 (0.21)	-1.34 (0.54)	-19.73 (0.65)	-0.86 (0.031)	2.78 (0.10)
<b>Cold Carcass Weight (kg)</b>	ns	2.12 (0.37)	-1.39 (0.44)	-0.19 (0.021)	-0.23 (0.070)
<b>Conformation (15 points)</b>	0.017 (0.006)	0.062 (0.015)	-0.043 (0.018)	-0.005 (0.001)	0.010 (0.003)
<b>Fat Class (15 points)</b>	0.072 (0.008)	0.199 (0.020)	-0.033 (0.024)	ns	-0.014 (0.001)
<b>Carcass Growth Rate (kg/day)</b>	-0.00603 (0.00025)	-0.00216 (0.00063)	0.00730 (0.00077)	0.00060 (0.00004)	-0.00142 (0.00012)
<b>Calf 200-day weight (kg)</b>	-7.19 (1.90)	18.11 (3.98)	-20.82 (7.14)	ns	ns
<b>Calf 200-day growth (kg/day)</b>	-0.0528 (0.0031)	-0.0332 (0.0058)	-0.0490 (0.0096)	0.0060 (0.0003)	-0.0083 (0.0021)

### 3.2 Extreme Weather

In the models including extreme weather frequencies, where effects were significant ( $p<0.05$ ) an increased frequency of extreme weather days had a negative effect on almost all traits (Table 2), assuming that a reduced age at slaughter and increased fat classes are desirable. Only for conformation score was an increase in frequency of dry days and wet days associated with improved conformation score. The effect of frequency of cold waves was only significant for conformation score ( $p<0.05$ ), where an increase in frequency of cold waves experienced was associated with a lower conformation score. For the calf traits, fewer types of extreme days had effects. Calf

200-day weight was only affected by the frequency of heatwaves and calf growth was only affected by the frequency of heatwaves and dry days.

**Table 2**

Table of solutions and standard errors (in brackets) for number of extreme weather days per day of life, from model for each carcass trait and calf trait. All effects are significant where given ( $p < 0.05$ ). Non-significant effects are denoted by ns.

	<b>Heatwaves</b>	<b>Cold Waves</b>	<b>Dry Days</b>	<b>Wet Days</b>
<b>Age at Slaughter (days)</b>	312.5 (10.7)	ns	167.3 (1.78)	83.18 (1.04)
<b>Cold Carcass Weight (kg)</b>	-20.44 (7.34)	ns	-13.51 (1.23)	-16.84 (0.72)
<b>Conformation (15 points)</b>	-0.80 (0.30)	-4.25 (0.78)	0.31 (0.051)	0.26 (0.030)
<b>Fat Class (15 points)</b>	-2.61 (0.40)	ns	-0.61 (0.067)	-0.53 (0.039)
<b>Carcass Growth Rate (kg/day)</b>	-0.18 (0.013)	ns	-0.082 (0.0022)	-0.050 (0.0013)
<b>Calf 200-day weight (kg)</b>	-1.29 (0.57)	ns	ns	ns
<b>Calf 200-day growth (kg/day)</b>	-0.010 (0.0026)	ns	-0.0065 (0.0011)	ns

#### 4. Conclusions

In conclusion, our results show that varying weather and frequency of extreme weather, across the lifetime of a beef animal, influences traits which affect the potential profit for a beef farmer. For example, we predict a 1°C increase in average daily maximum temperatures would reduce carcass growth rates by about 6g per day and calf growth rates by about 50g per day. We also predict an increase in frequency of heatwaves by 1 heatwave day per 100 days of life would reduce cold carcass weights by about 200g and increase age at slaughter by about 3 days.

These effects may be due to several factors, including direct effects on the animal, as well as feed availability and management decisions made by the farmer. However, there is potential to mitigate negative effects through a range of strategies, including selective breeding of animals with improved resilience to weather.

#### 5. Partners involved in the work

SRUC

#### 6. Annex – Paper

Paper submitted to animal: The Effects of Weather on Beef Carcass and Growth Traits. H. Bunning, E. Wall.

GenTORE – GA n° 727213

D6.2 - Paper on identification of key weather perturbations in performance and effect of decrease in temperature

1   **The Effects of Weather on Beef Carcass and Growth Traits**

2   H. Bunning<sup>a</sup>, E. Wall<sup>a</sup>

3   <sup>a</sup>*Animal Veterinary Science, SRUC, Edinburgh, UK, EH25 9RG*

4   Corresponding author: H. Bunning, Email: [harriet.bunning@sruc.ac.uk](mailto:harriet.bunning@sruc.ac.uk)

5   **Abstract**

6   To predict the effects of climate change, we need a better understanding of how beef  
7   cattle traits are affected by varying weather and frequency of extreme events. We  
8   analysed the effect of average daily precipitation and maximum and minimum  
9   temperatures on a range of important carcass traits, including age at slaughter, cold  
10   carcass weight, carcass growth rate and conformation and fat score ( $N = >1.6$   
11   million), as well as calf 200-day live weight and growth rate ( $N = >270\,000$ ), using  
12   data abattoirs across Scotland, England and Wales and calves in herds across  
13   Scotland. Animals which experienced higher daily maximum and lower daily  
14   minimum temperatures had poorer age at slaughter and carcass growth rates and  
15   calves had poorer 200-day weights. Increased precipitation also led to poorer cold  
16   carcass weights, conformation scores, calf 200-day weights and calf growth. We also  
17   analysed the effect of frequency of extreme weather events, including heatwaves,  
18   cold waves, and dry and wet days. The frequency of heatwaves, dry and wet days  
19   were shown to have significant negative effects on almost all traits considered, for  
20   example, predicting that an increase in frequency of heatwaves by 1 day per 100  
21   days of life would reduce cold carcass weights by about 200g and increase age at  
22   slaughter by about 3 days. Results show that varying weather and frequency of  
23   extreme weather, across the lifetime of a beef animal, influences traits which affect

24 the potential profit for a beef farmer. These effects may be due to several factors,  
25 including direct effects on the animal, as well as feed availability and management  
26 decisions made by the farmer. However, there is potential to mitigate negative  
27 effects through a range of strategies.

28 **Keywords:** Climate, Cattle, Heat stress, Extreme weather, Resilience

29 **Implications**

30 Our results show that varying weather and frequency of extreme weather events  
31 experienced by a beef animal, influences important beef traits. We predict a 1°C  
32 increase in average daily maximum temperatures would reduce carcass growth rates  
33 by about 6g per day and calf growth rates by about 50g per day. We also predict an  
34 increase in frequency of heatwaves by 1 heatwave day per 100 days of life would  
35 reduce cold carcass weights by about 200g and increase age at slaughter by about 3  
36 days. Without mitigation, these effects could reduce profit for farmers as well as  
37 increasing environmental impact.

38 **Introduction**

39 Climate change predictions show UK weather is likely to change significantly over  
40 the coming decades, both in terms of average weather conditions but also the  
41 frequency of extreme weather events (European Economic Area, 2017). There is a  
42 need for British livestock farming to adapt to these challenges, both to maintain  
43 profits for farmers, but also to reduce further climate and environmental impacts  
44 (Wreford and Topp, 2020). However, to plan potential mitigation strategies, we need  
45 to understand how varying climate impacts UK livestock farming.

46 There is good evidence that cattle are affected by climate. In the tropics, cattle  
47 experiencing high temperatures (especially combined with high humidity) experience  
48 heat stress which has negative impacts on milk production (Mbuthia et al., 2021),  
49 health and fertility (Polsky and von Keyserlingk, 2017; Bagath et al., 2019; Herbut et  
50 al., 2019) and growth (Brown-Brandl, 2018). Despite cattle in the UK not  
51 experiencing these same high temperatures, studies show that even Scottish dairy  
52 cattle experience a drop in milk yield due to both extreme highs and lows in  
53 temperature (Hill and Wall, 2014). Cold weather also effects other cattle traits.  
54 Studies have shown animals that are more exposed to cold weather during winter  
55 have lower growth rates (Holmes et al., 1984) and the use of calf jackets, particularly  
56 for dairy calves is thought to mitigate this (Robertson, 2020). We also expect  
57 precipitation to have an effect on cattle traits, as it affects plant growth (Dellar et al.,  
58 2018) and will likely affect grazing feed quality and availability.

59 The majority of these studies consider the effects on dairy cattle and we lack large  
60 scale studies on the effects of weather in temperate environments on beef cattle.  
61 Beef cattle have higher upper critical temperatures than dairy cattle (Wreford and  
62 Topp, 2020), so may be less affected by heat stress. However typically UK beef  
63 cattle are not housed as much as dairy (Smith et al., 2001) which may mean they are  
64 more greatly affected by weather. Therefore our aim is to investigate the effect of  
65 weather and frequency of extreme weather events on a range of cattle traits  
66 important to beef production.

## 67 **Materials and methods**

68 To investigate the effects of weather on beef production, we analysed two datasets,  
69 one consisting of slaughter records from UK abattoir companies across England,

70 Wales and Scotland (Pritchard et al., 2021) and one consisting of calf records in  
71 Scotland recorded through the Scottish Government's Beef Efficiency Scheme.  
72 These were both combined with data supplied from British Cattle Movement Service  
73 (BCMS) and weather data from the MetOffice HadUK-Grid database (Perry, 2004;  
74 Hollis and Perry, 2005). Final datasets after removing animals with missing  
75 information contained over 1 680 000 abattoir records from animals alive between  
76 2000 to 2019 and over 270 000 calf records from calves alive between 2016-2019.

77 ***Animal parameters***

78 Carcass traits included cold carcass weight (CCW), conformation class and fat class.  
79 Typically, conformation is assessed using the EUROP classification and fat class  
80 using a 1-5 scale. However, most abattoirs further sub-divide these classes.  
81 Therefore, these data were transformed to two 15 point scales, where 15 represents  
82 the best conformation and the fattest carcasses. Age at slaughter (AAS) was  
83 calculated using date of birth from BCMS data and kill date from abattoir data. A  
84 measure of carcass growth rate was calculated by dividing CCW by AAS. We call  
85 this average daily carcass gain (ADCG), but it is important to note that we have  
86 omitted birth weight in this calculation for simplicity as birth weight data was  
87 unavailable. Edits were made to remove extreme records, including those more than  
88 3 standard deviations from the mean of CCW, animals which were less than 365 or  
89 more than 1095 days old at slaughter and those with an ADCG more than 3 standard  
90 deviations from the mean. As well carcass data from abattoirs, we also had live  
91 weights for over 270 000 calves in Scotland, measured at approximately 200 days.  
92 The actual age at weighing varied from 100-300 days. We used these values to also  
93 calculate a calf growth rate trait, dividing the live weight by age at weighing.

94 A range of other factors and covariates were included. Sex was defined using data  
95 from the abattoir as castrated male (n=934,341 or 56%), female (n=527,741 or 31%)  
96 or entire male (n=219,722 or 13%). This was checked using data from BCMS where  
97 animals were recorded as male or female. For calves, we did not have information  
98 about castration status so all calves were recorded as male or female. Breed was  
99 defined using the breed code recorded in BCMS. Only animals from breeds with  
100 more than 1000 animals were included, resulting in the inclusion of 47 breeds. The  
101 most common three were Aberdeen Angus cross (n=287,687), Limousin cross  
102 (273,081) and Holstein (212,256).

103 Data about the dam of each animal was also extracted from BCMS. This included  
104 the age of the dam at the birth of the animal. Only individuals with dams older than  
105 365 days were included. This resulted in a dam age range of 371-3649 days with a  
106 mean of 1787 days and a standard deviation of 752 days. We also included the  
107 proportion of dairy breed in the dam's pedigree as this has been shown to have an  
108 important effect on carcass traits, particularly conformation score (Pritchard et al.,  
109 2021). This varied from 0.03-100%, with a mean of 80.22% and a standard deviation  
110 of 28.31%.

111 We needed to account for varying management practices which might be regionally  
112 distributed and therefore correlated with weather. We achieved this by including two  
113 contemporary groups in our model. First, we grouped animals according to their birth  
114 location, year and season (BirthHYS), where season was defined as three classes  
115 (Feb - May; Jun - Sep; Oct – Jan). We only included animals in BirthHYS groups that  
116 contained at least 5 animals. For the abattoir data, this resulted in 111,895 BirthHYS  
117 groups, ranging in size from 5 to 527, with a mean size of 15.0 animals. Secondly for  
118 the abattoir data only, we grouped animals according to their finishing location, year

119 and season (FinishHYS). We defined finished location as the location where an  
120 animals stayed for at least 60 days before slaughter (excluding up to 7 days before  
121 death to account for holding animals were moved through before slaughter). This  
122 resulted in 53,994 FinishHYS groups, ranging in size from 5 to 975 animals with a  
123 mean size of 31.2 animals. Finally, for the abattoir data, the location of death was  
124 also included. There were 32 death locations with between 785 and 181,494 animal  
125 slaughter records.

126 ***Weather parameters***

127 We used weather data from the Met Office HadGrid-UK database, a data set of  
128 gridded climate variables derived from the network of UK land surface observations.  
129 Variables include daily maximum and daily minimum temperatures and daily total  
130 precipitation for each 1km square across the UK. Animal locations and dates of stay  
131 were extracted from the BCMS database and the nearest centre of a corresponding  
132 km square from the HadGrid data found. This allowed us to calculate the average  
133 daily maximum temperature (Tmax), average daily minimum temperature (Tmin) and  
134 average daily precipitation (Rain) for the lifetime of each animal.

135 The daily weather was also used to define the occurrence of extreme weather  
136 events, including heatwaves, coldwaves, dry days and wet days. The Met Office  
137 definition of a heatwave is a period of at least 3 days where the daily maximum  
138 temperature exceeds a threshold. The threshold is specific to the location, with four  
139 threshold regions defined by the met office in the UK: London, the South East of  
140 England, an area around the South East of England and the rest of the UK, with  
141 thresholds of 28°C, 27°C, 26°C and 25°C respectively. For cold waves, a similar  
142 definition was used, where a period consisted of at least 3 days where the daily

143 maximum temperature did not exceed 0°C. Wet and dry days were defined as days  
144 where rainfall was greater than 7.65mm and less than 0.12mm respectively. These  
145 values correspond to 90<sup>th</sup> and 10<sup>th</sup> percentile of the daily precipitation across the UK  
146 for the period 2000-2019. For wet and dry days no minimum number of consecutive  
147 days was required. The total number of each type of extreme day experienced by  
148 each animal was calculated and divided by its AAS or age at weighing for calves, to  
149 calculate the frequencies of extreme weather days.

150 ***Statistical analysis of results***

151 Analyses were carried out using linear fixed effect models using AS-REML and R.  
152 Two models were produced for each trait, the first to assess the effect of average  
153 weather and the second to assess the effect frequency of extreme weather events.  
154 For each carcass trait, all other carcass traits, except ADCG, were included as  
155 covariates. For ADCG, AAS and CCW were also not included. For the two calf traits  
156 (calf weight and calf growth), no other traits and no FinishHYS or death location were  
157 included. We expected interactions between weather to be important so an  
158 interaction between Tmax and Tmin and another between Tmin and Rain were  
159 included in the average weather models. The generalised model was therefore as  
160 follows:

161 *Trait ~ weather parameters + other traits + sex + breed + BirthHYS + FinishHYS +*  
162 *death location + dam age + dam %dairy*

163 **Results**

164 **Average weather**

165 Almost all average weather parameters had a significant ( $p<0.05$ ) effect on every  
166 trait assessed (Table 1). An increase in AAS, which is undesirable as increases  
167 farmer costs, was seen in animals which experienced higher Tmax ( $\beta = 10.17$ , s.e. =  
168 0.21), lower Tmin ( $\beta = -1.34$ , s.e. = 0.54) and lower Rain ( $\beta = -19.73$ , s.e. = 0.65).  
169 The effect of the interactions between Tmin-Tmax ( $\beta = -0.86$ , s.e. = 0.031) and Tmin-  
170 Rain ( $\beta = 2.78$ , s.e. = 0.10) were also significant for AAS. CCW was not significantly  
171 affected by Tmax ( $p > 0.05$ ), but higher weights were associated with higher Tmin ( $\beta$   
172 = 2.12, s.e. = 0.37) and lower Rain ( $\beta = -1.39$ , s.e. = 0.44). Again, the effect of the  
173 interactions between Tmin-Tmax ( $\beta = -0.19$ , s.e. = 0.021) and Tmin-Rain ( $\beta = -0.23$ ,  
174 s.e. = 0.070) were also significant. Higher conformation scores were seen for  
175 animals which experienced high Tmax ( $\beta = 0.017$ , s.e. = 0.006) and Tmin ( $\beta = 0.062$ ,  
176 s.e. = 0.015) and lower Rain ( $\beta = -0.043$ , s.e. = 0.018). Interactions between Tmin-  
177 Tmax ( $\beta = -0.005$ , s.e. = 0.001) and Tmin-Rain ( $\beta = 0.010$ , s.e. = 0.003) were also  
178 shown to have a significant effect on conformation score. An increase in fat score  
179 was seen in animals which experienced higher Tmax ( $\beta = 0.072$ , s.e. = 0.008) and  
180 Tmin ( $\beta = 0.199$ , s.e. = 0.020) and lower Rain ( $\beta = -0.033$ , s.e. = 0.024). For fat  
181 score, only the interaction between Tmin and Rain ( $\beta = -0.014$ , s.e. = 0.001) was  
182 significant ( $p<0.05$ ). For ADCG, higher growth rates were associated with animals  
183 that experiences lower Tmax ( $\beta = -0.0060$ , s.e. = 0.00025) and Tmin ( $\beta = -0.0022$ ,  
184 s.e. = 0.00063) and higher Rain ( $\beta = 0.0073$ , s.e. = 0.00077). Again, interactions  
185 between Tmin-Tmax ( $\beta = 0.00060$ , s.e. = 0.00004) and Tmin-Rain ( $\beta = -0.0014$ , s.e.  
186 = 0.00012) were also shown to have a significant effect on ADCG.

187 For the calf traits, greater 200-day live weights were associated with animals that  
188 had experienced lower Tmax ( $\beta = -7.19$ , s.e. = 1.90), higher Tmin ( $\beta = 18.11$ , s.e. =  
189 3.98) and lower Rain ( $\beta = -20.82$ , s.e. = 7.14). Interactions between weather effects

190 were not significant ( $p>0.05$ ). An increase in calf growth rate was seen for animals  
191 that had experienced lower Tmax ( $\beta = -0.053$ , s.e. = 0.0031), Tmin ( $\beta = -0.033$ , s.e.  
192 = 0.0058) and Rain ( $\beta = -0.049$ , s.e. = 0.0096). Interactions between Tmin-Tmax ( $\beta =$   
193 0.0060, s.e. = 0.0003) and Tmin-Rain ( $\beta = -0.0083$ , s.e. = 0.0021) were also shown  
194 to have a significant effect on calf growth rate.

195 ***Extreme weather***

196 In the models including extreme weather frequencies, where effects were significant  
197 ( $p<0.05$ ) an increased frequency of extreme weather days had a negative effect on  
198 almost all traits (Table 2), assuming that a reduced AAS and increased fat classes  
199 are desirable. Only for conformation score was an increase in frequency of dry days  
200 ( $\beta = 0.31$ , s.e. = 0.051) and wet days ( $\beta = 0.26$ , s.e. = 0.030) associated with  
201 improved conformation score. The effect of frequency of cold waves was only  
202 significant for conformation score ( $p<0.05$ ), where an increase in frequency of cold  
203 waves experienced was associated with a lower conformation score ( $\beta = -4.25$ , s.e.  
204 = 0.78). For the calf traits, fewer types of extreme days had effects. Calf 200-day  
205 weight was only affected by the frequency of heatwaves ( $\beta = -1.29$ , s.e. = 0.57) and  
206 calf growth was only affected by the frequency of heatwaves ( $\beta = -0.010$ , s.e. =  
207 0.0026) and dry days ( $\beta = -0.0065$ , s.e. = 0.0011).

208 **Discussion**

209 It is clear from these results that varying weather across the lifetime of a beef animal  
210 influences traits which affect the potential profit for a beef farmer. These effects may  
211 be due to several factors, including the effects of weather on feed quality and  
212 availability, management decisions made by the farmer and the physiology and  
213 behaviour of the animal (Wreford and Topp, 2020).

214 An increase in average daily maximum temperature led to poorer AAS, calf weight  
215 and calf and carcass growth rates, but improved conformation and higher fat class.  
216 Animals which have experienced high average temperatures (especially alongside  
217 high humidity which we were unable to account for in these analyses) are more likely  
218 to have experienced heat stress, which has been shown to have a detrimental  
219 impact on growth rate in beef cattle, due to both reduced feed intake but also direct  
220 effects on metabolism (Brown-Brandl, 2018). Typically, these effects are considered  
221 in countries with warmer climates, but effects have been seen in UK dairy cattle  
222 where extremes of THI led to reduced milk yields (Hill and Wall, 2014). The threshold  
223 where UK animals will be affected will be much lower than those acclimated to  
224 warmer climates (Collier et al., 2019) which is why we expect to see effects even at  
225 the lower temperatures seen in the UK. Supporting this, our results show that  
226 animals which experience an increased number of heat waves days per day of life  
227 tend to have poorer AAS, conformation and fat score and carcass and calf weights  
228 and growth rates. On these extreme hot days, cattle feed less, both to avoid leaving  
229 shaded areas but also to reduce heat production in the rumen, as well as expending  
230 additional energy to attempt to dissipate heat (Van Laer et al., 2014).

231 An increase in the average daily minimum temperature experienced by an animal  
232 has some similar effects to those seen for maximum temperature. However, whereas  
233 calf weights were reduced and CCW was not significantly affected with increasing  
234 maximum temperatures, both carcass and calf weights increased with increasing  
235 minimum temperatures. Cold temperatures will reduce forage yields as growth is  
236 limited (Hurtado-Uria et al., 2013), which may lead to reduced weights if feed  
237 availability is also limited, but this may be mitigated by supplementary feeding. Cold  
238 temperatures will also have a direct impact on the physiology of the animal. Outside

239 the boundaries of the thermo-neutral zone, animals must expend energy, in this case  
240 to remain warm (Van laer et al., 2014). This lower limit is higher for calves than adult  
241 animals (Van laer et al., 2014) so we expect their weights to be more negatively  
242 affected, which is in line with our results. One unexpected result is the increase in fat  
243 class seen under increasing daily minimum temperatures. We might expect animals  
244 experiencing less cold weather to have reduced levels of subcutaneous fat,  
245 decreasing the fat score (Van laer et al., 2014). Our result may be due to the  
246 reduced energy requirements for maintenance under warmer daily minimum  
247 temperatures, allowing more energy to be stored as fat. Despite the important effects  
248 of average daily minimum temperature, we did not see significant effects for  
249 frequency of cold waves, except for a decrease in conformation score (which is in  
250 line with the effect of average daily minimum temperature). This is possibly due to  
251 the relatively number of cold waves seen within the dataset compared to heatwaves.

252 Our results show that increased rainfall leads to a poorer CCW, conformation score,  
253 fat score, calf weight and calf growth rate. Increased rainfall is associated with  
254 increased risk of fluke infection (Skuce et al., 2014). Presence of a fluke infection  
255 has been shown to be associated with reduced CCW and lower conformation and fat  
256 scores (Bellet et al., 2016) which corresponds with our results. However increased  
257 rainfall also led to improved growth rates for abattoir animals and lower AAS. This  
258 beneficial effect seems unlikely to be due to a direct effect on either the physiology  
259 or behaviour of the cattle, therefore this is more likely to be due to either a change in  
260 feed availability or some other change in management. Indeed, we expect increased  
261 rainfall to lead to improved pasture yields (Dellar et al., 2018) which could account  
262 for this increase in growth rate and reduced age to slaughter. However, when we  
263 consider the number of extreme wet days experienced by an animal, we predict a

264 reduction in carcass growth rates and poorer AAS, showing that although generally  
265 more rain may have some beneficial effects, days of extreme wet weather are  
266 detrimental to growth. This could be due to several factors, including a change in  
267 animal behaviour during these extreme periods which leads to reduced feeding  
268 either to avoid rain or even flooding. Alternatively, these could reflect damage to  
269 pastures leading to reduced feed availability or changes in management surrounding  
270 these days, for example limited access to provide supplementary feed. Extreme dry  
271 days also led to poorer AAS, CCW and both carcass and calf growth rates. This is  
272 unlikely to be a direct effect on the physiology of animal, as animals will have water  
273 provisions even during dry periods. The effect is more likely due to a reduced  
274 pasture yield and quality as grass growth is severely limited during dry periods  
275 (Dellar et al., 2018). This reduces feed quality and availability for grazing animals.

276 Current climate change projections suggest that in the UK summer and winter  
277 temperatures will increase, whilst summer rainfall will decrease and winter rainfall  
278 will increase (Wreford and Topp, 2020). Without changes to management or  
279 acclimatisation of cattle, these changes may lead to some negative impacts to beef  
280 production. We predict a 1°C increase in average daily maximum temperatures  
281 would reduce carcass growth rates by about 6g per day and calf growth rates by  
282 about 50g per day. These effects could lead to longer time to slaughter or reduced  
283 carcass weights, reducing the potential profit for farmers as well as increasing  
284 environmental impact by increasing GHG emissions.

285 Unlike the more gradual change in climate, animals are unlikely to acclimatise to  
286 extreme weather events (Collier et al., 2019) and these may also be more difficult to  
287 mitigate through management changes. Frequency of these extreme events are  
288 likely to increase (European Economic Area, 2017) and our results predict a negative

289 impact of this on almost all traits. For example, our results predict an increase in  
290 frequency of heatwaves by one heatwave day per 100 days of life would reduce  
291 CCW by about 200g and increase AAS by about three days, again reducing the  
292 potential profit for farmers as well as increasing environmental impact.

293 There is potential to reduce these effects through a number of varying strategies.  
294 Planting more hedges and trees around pastures to provide cover could negate the  
295 negative effects of heat, cold and rain on the animal (Van laer et al., 2014) and this  
296 strategy would be relatively inexpensive and potentially provide environmental  
297 benefits (Forman and Baudry, 1984). More substantial shelter could also be provided  
298 in the form of housing, particularly for some outwintered cattle. For housed cattle  
299 experiencing heat stress, better ventilation could be used to mitigate the negative  
300 effects (Van laer et al., 2014). Where weather affects pasture growth, more  
301 supplementary feeding may be required, although this may be costly, both for farmer  
302 profit but also environmental impact (Sasu-Boakye et al., 2014). In addition to these  
303 strategies, farmers may want to consider selecting breeds or genotypes which are  
304 more resilient and therefore less affected by varying weather (Sánchez-Molano et  
305 al., 2020; Poppe et al., 2021).

306 In conclusion, our results show that that varying weather and frequency of extreme  
307 weather, across the lifetime of a beef animal, influences traits which affect the  
308 potential profit for a beef farmer. These effects may be due to several factors,  
309 including direct effects on the animal, as well as feed availability and management  
310 decisions made by the farmer. However, there is potential to mitigate negative  
311 effects through a range of strategies.

312 **Ethics approval**

313 Not applicable

314 **Author contributions**

315 **H Bunning:** Methodology, Formal analysis, Data curation, Writing **E Wall:**

316 Conceptualization, Supervision, Project administration, Funding acquisition

317 **Declaration of interest**

318 None

319 **Acknowledgements**

320 None

321 **Financial support statement**

322 The research leading to these results has received funding from European Union's

323 Horizon 2020 research and innovation programme - GenTORE - under grant

324 agreement N° 727213.

325 Research and data collection as part of the Beef Efficiency Scheme funded by

326 Scottish Government Strategic Research Programme 2016-2021.

327 **References**

328 Bagath M, Krishnan G, Devaraj C, Rashamol VP, Pragna P, Lees AM and Sejian V

329 2019. The impact of heat stress on the immune system in dairy cattle: A review.

330 Research in Veterinary Science 126, 94–102.

331 Bellet C, Green MJ, Vickers M, Forbes A, Berry E and Kaler J 2016. Ostertagia spp.,

332 rumen fluke and liver fluke single- and poly-infections in cattle: An abattoir study of

333 prevalence and production impacts in England and Wales. Preventive Veterinary

- 334 Medicine 132, 98–106.
- 335 Brown-Brandl TM 2018. Understanding heat stress in beef cattle. Revista Brasileira  
336 de Zootecnia 47.
- 337 Collier RJ, Baumgard LH, Zimbelman RB and Xiao Y 2019. Heat stress: Physiology  
338 of acclimation and adaptation. Animal Frontiers 9, 12–19.
- 339 Dellar M, Topp CFE, Banos G and Wall E 2018. A meta-analysis on the effects of  
340 climate change on the yield and quality of European pastures. Agriculture,  
341 Ecosystems and Environment 265, 413–420.
- 342 European Economic Area 2017. Climate change, impacts and vulnerability in Europe  
343 2016. Copenhagen, Denmark.
- 344 Forman RTT and Baudry J 1984. Hedgerows and hedgerow networks in landscape  
345 ecology. Environmental Management 1984 8:6 8, 495–510.
- 346 Herbut P, Angrecka S, Godyń D and Hoffmann G 2019. The Physiological and  
347 Productivity Effects of Heat Stress in Cattle-A Review. Annals of Animal Science 19,  
348 579–593.
- 349 Hill DL and Wall E 2014. Dairy cattle in a temperate climate: The effects of weather  
350 on milk yield and composition depend on management. Animal 9, 138–149.
- 351 Hollis D and Perry M 2005. Development of a new set of long-term climate averages  
352 for the UK. International Journal of Climatology 25, 1023–1039.
- 353 Holmes CW, Christensen R, McLean NA and Lockyer J 1984. Effects of winter  
354 weather on the growth rate and heat production of dairy cattle. New Zealand Journal  
355 of Agricultural Research 21, 549–556.

- 356 Hurtado-Uria C, Hennessy D, Shalloo L, O'Connor D and Delaby L 2013.
- 357 Relationships between meteorological data and grass growth over time in the south  
358 of Ireland. *Irish Geography* 46, 175–201.
- 359 Van laer E, Moens CPH, Sonck B and Tuyttens FAM 2014. Importance of outdoor  
360 shelter for cattle in temperate climates. *Livestock Science* 159, 87–101.
- 361 Mbuthia JM, Mayer M and Reinsch N 2021. Modeling heat stress effects on dairy  
362 cattle milk production in a tropical environment using test-day records and random  
363 regression models. *Animal* 15, 100222.
- 364 Perry M 2004. The generation of monthly gridded datasets for a range of climatic  
365 variables over the United Kingdom. Met Office, Exeter, UK, 1–20.
- 366 Polsky L and von Keyserlingk MAG 2017. Invited review: Effects of heat stress on  
367 dairy cattle welfare. *Journal of Dairy Science* 100, 8645–8657.
- 368 Poppe M, Mulder HA and Veerkamp RF 2021. Validation of resilience indicators by  
369 estimating genetic correlations among daughter groups and with yield responses to a  
370 heat wave and disturbances at herd level. *Journal of Dairy Science* 104, 8094–8106.
- 371 Pritchard TC, Wall E and Coffey MP 2021. Genetic parameters for carcase  
372 measurements and age at slaughter in commercial cattle. *Animal* 15, 100090.
- 373 Robertson J 2020. Calf jackets: a review of science and practice. *Livestock* 25, 284–  
374 290.
- 375 Sánchez-Molano E, Kapsona V V., Oikonomou S, McLaren A, Lambe N, Conington  
376 J and Banos G 2020. Breeding strategies for animal resilience to weather variation in  
377 meat sheep. *BMC Genetics* 21, 1–11.

- 378 Sasu-Boakye Y, Cederberg C and Wirsénus S 2014. Localising livestock protein  
379 feed production and the impact on land use and greenhouse gas emissions. animal  
380 8, 1339–1348.
- 381 Skuce P, Van Dijk J, Smith D and Morgan E 2014. Editorial: Implications of extreme  
382 weather events for risk of fluke infection. Veterinary Record 175, 198–200.
- 383 Smith KA, Brewer AJ, Crabb J and Dauven A 2001. A survey of the production and  
384 use of animal manures in England and Wales. III. Cattle manures. Soil Use and  
385 Management 17, 77–87.
- 386 Wreford A and Topp CFE 2020. Impacts of climate change on livestock and possible  
387 adaptations: A case study of the United Kingdom. Agricultural Systems 178, 102737.
- 388
- 389

390 **Tables**391 **Table 1**

392 Table of solutions and standard errors (in brackets) for weather variables and  
 393 interactions from model for each carcass trait and calf trait. All effects are significant  
 394 where given ( $p<0.05$ ). Non-significant effects are denoted by ns.

	Tmax	Tmin	Rain	Tmin x Tmax	Tmin x Rain
<b>Age at Slaughter (days)</b>	10.17 (0.21)	-1.34 (0.54)	-19.73 (0.65)	-0.86 (0.031)	2.78 (0.10)
<b>Cold Carcass Weight (kg)</b>	ns	2.12 (0.37)	-1.39 (0.44)	-0.19 (0.021)	-0.23 (0.070)
<b>Conformation (15 points)</b>	0.017 (0.006)	0.062 (0.015)	-0.043 (0.018)	-0.005 (0.001)	0.010 (0.003)
<b>Fat Class (15 points)</b>	0.072 (0.008)	0.199 (0.020)	-0.033 (0.024)	ns	-0.014 (0.001)
<b>ADCG (kg/day)</b>	-0.00603 (0.00025)	-0.00216 (0.00063)	0.00730 (0.00077)	0.00060 (0.00004)	-0.00142 (0.00012)
<b>Calf 200-day weight (kg)</b>	-7.19 (1.90)	18.11 (3.98)	-20.82 (7.14)	ns	ns
<b>Calf 200-day growth (kg/day)</b>	-0.0528 (0.0031)	-0.0332 (0.0058)	-0.0490 (0.0096)	0.0060 (0.0003)	-0.0083 (0.0021)

395

396

397 **Table 2**

398 Table of solutions and standard errors (in brackets) for number of extreme weather  
 399 days per day of life, from model for each carcass trait and calf trait. All effects are  
 400 significant where given ( $p<0.05$ ). Non-significant effects are denoted by ns.

	<b>Heatwaves</b>	<b>Cold Waves</b>	<b>Dry Days</b>	<b>Wet Days</b>
<b>Age at Slaughter (days)</b>	312.5 (10.7)	ns	167.3 (1.78)	83.18 (1.04)
<b>Cold Carcass Weight (kg)</b>	-20.44 (7.34)	ns	-13.51 (1.23)	-16.84 (0.72)
<b>Conformation (15 points)</b>	-0.80 (0.30)	-4.25 (0.78)	0.31 (0.051)	0.26 (0.030)
<b>Fat Class (15 points)</b>	-2.61 (0.40)	ns	-0.61 (0.067)	-0.53 (0.039)
<b>ADCG (kg/day)</b>	-0.18 (0.013)	ns	-0.082 (0.0022)	-0.050 (0.0013)
<b>Calf 200-day weight (kg)</b>	-1.29 (0.57)	ns	ns	ns
<b>Calf 200-day growth (kg/day)</b>	-0.010 (0.0026)	ns	-0.0065 (0.0011)	ns

401